Host Preference, Mode of Damage and Succession of Major Insect Pests of Brinjal

Saifullah Omar Nasif¹ and Shafiquzzaman Siddiquee^{2*}

¹*Department of Entomology, Sher-e-Bangla Agricultural University, Dhaka, Bangladesh.*

²*Biotechnology Research Institute, Universiti Malaysia Sabah, Jln UMS, 88400 Kota Kinabalu, Sabah, Malaysia.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2020/v35i830261

Editor(s):

(1) Dr. Bechan Sharma, University of Allahabad, India.

Reviewers:

(1) Surajit Kalita, Assam Agricultural University (AAU), India.

(2) Uzma Manzoor, Sharda University, India.

(3) Md. Ramjan, College of Horticulture and Forestry, CAU, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/59627>

Review Article

Received 22 May 2020
Accepted 27 July 2020
Published 13 August 2020

ABSTRACT

Brinjal is one of the most popular solanaceous vegetables worldwide because of its nutritive and nutraceutical values. In recent decades, brinjal cultivation has been experiencing increased biotic and abiotic stresses due to the climate changes, which had affected the cultivation practices, growers' profit and consumers' choice to a larger extent. Amongst different insect pests, brinjal shoot and fruit borer (BSFB) is the major one causing significant reduction in economic yield followed by epilachna beetle, aphid, jassid and whitefly. On the other hand, proper knowledge on nature of damage, host preference, and seasonal abundance of insect pests is a pre-requisite for development and deployment of an appropriate, effective and timely management strategy against the pests. A pool of literatures on host preferences, damage pattern and succession of major insect pests of brinjal has been reviewed and presented in this paper.

Keywords: *Brinjal; insect pests; nutritive and nutraceutical values; vegetables.*

**Corresponding author: E-mail: shafiqpab@ums.edu.my;*

1. INTRODUCTION

Brinjal or eggplant (*Solanum melongena* L.) (Family: Solanaceae) is the most admired and predominant non-tuberous vegetable in Bangladesh and other parts of the world. The genus *Solanum* includes diverse flowering plants, including a few high-value economically important food crops like brinjal [1]. Global brinjal production is 51.23 m MT and China is the world's largest brinjal producer (32 m MT) followed by India (12.55 m MT) and Egypt (1.19 m MT) [2].

Brinjal is well known for its high-water content and low calorific value [3]. A brinjal fruit contains moisture 91.5 per cent along with 1.3 per cent protein, 6.5 per cent minerals, 6.4 per cent carbohydrates, 0.02 per cent calcium, 0.06 per cent phosphorus and 1.3 per cent iron, respectively [4]. It also a good source of vitamin A (5 mg/100 g), vitamin B (45 mg/ 100 g), nicotinic acid (0.08 mg/ 100 g), riboflavin (90 mg/ 100 g) and vitamin C (23 mg/ 100) [4]. Moreover, brinjal has been ranked to be amongst the top ten vegetables in terms of antioxidant capacity due to the presence of antioxidants such as arginine, 5-HT, delphinidine-3 Bioside (nasunin), solasodine, tryptophan, etc. [5,6]. Beside its food value, brinjal has been known for its medicinal values because of presence of phenols on fruits such as anthocyanins and strychnine, which have the potential to cure a variety of disease like cancer, hypertension, hepatosis [7,8]. Brinjal has also beneficial effects in the treatment of inflammatory stress, cardiac debility, neuralgias, bronchitis and asthma [9] and visual function [10] [11]. Also revealed that brinjal contains fraction of crude alkaloid that has significant analgesic effect.

Among the various causes of low productivity of the brinjal, damage inflicted by the insect pests is the most important one [12] causing 28-85% reduction in yield [13]. Some of the major insect pests attacking brinjal are, Brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guen.); Whitefly, *Bemisia tabaci* (Genn.); Epilachna beetle, *Epilachna vigintioctopunctata*; leafhopper, *Amrasca biguttula biguttula* (Ishida); Mealy bug, *Coccidohystrix insolita* (Green); Thrips, *Thrips palmi*, red mite (*Tetranychus urticae*) etc. The losses caused by various pests were Brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee (Lepidoptera: Crambidae) is the key pest [14,15,16] incurring severe damage in nearly all the brinjal cultivating nations [17], more

particularly, south Asia i.e. India, Bangladesh as a whole [18]. Sucking insect pests such as jassid, aphid and whitefly are also considered to be the serious pests of brinjal in Bangladesh [19]. Increase in maximum and minimum temperature leads to increase the number of generation whereas lower temperature lengthen the duration of life cycle of insect pests. Thus, increased climate change effect makes it harder to control insect pests effectively.

Although, several eco-friendly management strategies have been evolved to manage these pests, but to develop and design economically feasible and effective pest management strategies, information on bio-ecology along with ecological footing is a pre-requisite. Similarly, influence of weather factors on pest population along with sufficient knowledge on seasonal activity of the pest is also necessary for adopting location specific suitable control measures in a particular region. Hence, this review aims at compiling and presenting the distribution, nature of damage and succession of major insect pests of brinjal.

2. MAJOR INSECT PESTS OF BRINJAL

About 53 species of insect pests are found to attack brinjal cultivation worldwide [20] of which, 20 insect pests (19 insects and 1 mite pest) were reported to cause serious damage in Bangladesh [21]. Among them, brinjal shoot and fruit borer, epilachna beetle, jassid, aphid and whitefly were described to be the major insect pests of brinjal through pest risk analysis [21]. A detail report on host preference, damage and succession of major insect pest of brinjal has been discussed below.

2.1 Brinjal Shoot and Fruit Borer (BSFB), *Leucinodes orbonalis* Guen.

BSFB is the most notorious pest of brinjal in Bangladesh and the genus *Leucinodes* reported to consists of three main species namely *L. orbonalis* Guen., *L. diaphana* Hamps and *L. apicalis* Hamps [22].

2.1.1 Host preferences

Besides brinjal, BSFB is found to attack other solanaceous crops along with their wild relatives [23]. Caterpillar of this insect is also found to feed on pea pods [22]. *Solanum nigrum* and *Solanum myriacanthum* potentially plays significant role as

alternative host of brinjal shoot and fruit borer [24,25].

2.1.2 Nature of damage

L. orbonalis is found to attack flowering, fruiting and vegetative developing stage on fruits/units, developing parts and inflorescence of brinjal [24]. Like other holometabolans members of the insect order lepidoptera, *L. orbonalis* also undergoes four growth stages: egg, larva, pupa and adult. The larval period is the longest (11-13 days), followed by pupal (10-11 days) and incubation period (5-6 days) [26]. Oviposition takes place at night or in the morning hours of the day and eggs are laid singly on the lower surface of the young leaves, green stems, flower buds, or calyxes of the fruits [24]. The oviposition rate of the insect is reported to be 80 to 253 numbers of eggs per female [27,28]. Eggs are flattened, elliptical with 0.5 mm in diameter and color is creamy-white but change to red before hatching [29]. The egg takes incubation period of 3-5 days in summer and 7-8 days in winter and hatch into dark white larvae. The larval period lasts for 12 - 15 days during summer and 14 - 22 days during winter season [30]. Larvae pass through at least five instars [31,32] and there are reports of existence of six larval instars [33].

The caterpillars of *L. orbonalis* bore into the developing points of young tender shoots which at last shrivel away. The fruiting beads droop down while the fruits indicate round about openings. The higher percent of the larvae was found in fruits taken after by shoots, flowers, flower buds and midrib of leaves [28]. Inside one hour in the wake of bring forth, *L. orbonalis* caterpillar drills into the closest delicate shoot, bloom, or fruit and immediately blocks the passageway opening (nourishing passage) with excreta [29]. Watching the drilling openings, the pervaded fruits can be distinguished without much of a stretch. Furthermore, the dull shaded excreta can be seen without much effort to the opening of pervaded fruits. Optional pervasions by specific microorganisms may create additional decay of the fruits [34] and make them unfit for human consumption. In young plants, caterpillars are accounted for exhausting inside petioles and midribs of extensive leaves [28,35,36] along these lines shriveling, drop off and shrink of the young shoots prompting delay on shoot development, decrease on yield and yield parameter. Larval infestation inside the fruit brings about pulverization of fruit tissue and in serious cases, spoiling the whole fruit [37]. Larval

nourishing in flower was uncommon, if happen, inability to shape fruit from harmed flowers [29].

2.1.3 Succession of brinjal shoot and fruit borer

L. orbonalis is dynamic during the time at places having moderate atmosphere yet its movement is antagonistically influenced by serious chilling [38]. BSFB pervasion on brinjal starts in August and achieved its crest in October and afterward began declining [38]. According to [39], a low infestation (18.66%) of borer was noted in the third week of May, which got severe (75.50% infestation) in the first week of August, and moderate (42.64% infestation) in the last week of September at the end of the crop growing season [40]. Found that the pest is most dynamic amid the late spring months, i.e., from May to August. It turns out to be less dynamic amid the winter months, especially in December and January [41]. Considered the occurrence and plenitude of BSFB in Allahabad, India and watched the most elevated rate on brinjal in December [42]. Discovered shoot and fruit damage in brinjal by BSFB was higher in May transplanted (spring) crops than that of July and September transplanted (fall) crops. The damage caused by insect change from season to season since direct temperature and high moisture support the populace develop of brinjal shoot and fruit borer [43,44]. Areas with hot and humid climate are conducive for population buildup and higher incidence of the pest. [42] Reported that summer season brinjal has more susceptibility than winter season brinjal and he found highest shoot infestation during mid-September while peak fruit infestation was reported during mid-November.

2.2 Epilachna Beetle, *Epilachna dodecastigma*

Among the coccinellids, Epilachna beetles belonging to the subfamily Epilachninae constitute one-sixth species and around 500 species have been found under the genus *Epilachna* [45]. This pest is widely distributed in South East Asia, Australia, China, India and many other countries.

2.2.1 Host preference

Epilachna beetles are phytophagous and found to attack a wide range of plants belonging to solanaceae, cucurbitaceae, fabaceae,

convolvulaceae as well as malvaceae family. Brinjal, tomato, potato, tobacco, melon, cucumber, gourds, pumpkin and many other important food crops are frequently being under attack of epilachna beetle [46]. Its presence has also been reported from medicinal plants and other naturally occurring solanaceous plant like *Solanum nigrum* and *Solanum torvum* [47,48].

2.2.2 Nature of damage

Infestation of epilachna beetle can significantly reduce yield by hampering crop growth [49]. Both adult and grub scrap off the epidermal tissue of leaves, flowers and fruits inflicting serious damage to the brinjal plant from seedling stage up to maturity [40,41,50]. [51] Reported 35-75 percent leaf injury due to the attack of epilachna beetle attack, which could go upto 80 per cent [52]. While, [53] reported 60 percent fruit damage caused by this notorious pest. Its importance increases to several folds as [54] reported that epilachna beetle had developed resistance against many commercial insecticides in recent times.

2.2.3 Succession of epilachna beetle

According to [55], maximum temperature and minimum temperature has significant positive correlation with population dynamics of the pest, while negatively correlated with rainfall and humidity. In contrast, [56] reported that highest number of epilachna found during third week of February and reaching to the least during April. However, its infestation starts from the initial crop growth period up to maturity [56]. Also reported positive correlation of relative humidity and rainfall interms of succession and population dynamics of the epilachna beetle [41]. Reported that the infestation of epilachna get started by the first week of November with an average population of 2.85 beetles per plant reaching maximum in the third week of February with the first peak at the third week of November. They also revealed a negative correlation between temperature and populace dynamics of the pest, however positive correlation among all other weather parameters. Research carried out by [57] revealed that highest population of epilachna beetle found on mid-August which get declined by the end of August and became zero during October. The maximum temperature, minimum temperature and relative humidity during peak period of infestation of the pest were reported to be $27.5 \pm 0.88^{\circ}\text{C}$, $19.58 \pm 0.49^{\circ}\text{C}$ and

$75.55 \pm 13.37\%$ respectively [58]. Reported that infestation of epilachna beetle starts from December with a peak during February and March. However, [40] stated Mid-September to be the highly infested months. [59] Found $24-31^{\circ}\text{C}$ and 58-75% relative humidity have conducive effect in population build-up of epilachna beetle.

2.3 Jassid, *Amrasca biguttula biguttula* Ishida

Jassid is a common sucking pest of brinjal around the world with wide range of host preference and capability to cause damage.

2.3.1 Host preference

Besides brinjal, jassid attacks cotton crop, which also harbours on various herbaceous plants and weeds of solanaceae, malvaceae and Cruciferae family [60]. [61] claimed jassid as the second major pest of brinjal after BSFB due to its high population intensity and damage severity [62]. Reported that brinjal is one of the most favourite host plants of *A. biguttula biguttula*. Many researchers had identified jassid to be a key pest of Brinjal [63,64]. [64] stated that oriental regions i.e. tropical and subtropical regions are suitable for jassid population due to the fact that the weather conditions prevailing in these regions are conducive for host-plant interaction.

2.3.2 Nature of damage

Jassid is phytophagous in nature and the damage percentage of jassid could extend up to 54 percent [65]. Jassid could cause devastating effect in solanaceous crops by hampering transportation through the phloem tissues of plant and possibly introducing a toxin inhibiting photosynthesis activity [66]. Most importantly, they don't cause much economic damage by reducing plant vigor by sucking cell sap only, but they affect rigorously on fruit yield by spreading virus disease as a vector [67]. According to [68], jassids prefers brinjal cultivars those have smooth textured leaves than having leaves having leathery texture or leathery texture with spines. [62] reported that the hair density and length of hair on lamina, midrib, and veins of brinjal had highly significant and negative correlation with the jassid population. Moreover, trichome density on the leaves play important role in the plant defense particularly among phytophagous insects including jassids [69].

2.3.3 Succession of jassid on brinjal

A study on population dynamics of jassid [70] revealed first appearance of brinjal jassid during 32nd SMW and continues up to 41st SMW Highest number of jassids (12.70 jassids/ leaf) was reported during 37th SMW [71]. Carried out an experiment on succession of brinjal pest complex and found that jassid population appears in the first week after transplanting of the crop and continues up to the maturity [72]. Also reported the peak population of jassid in brinjal during 16th, 18th, 24th, 33rd SMW [73]. Observed peak population of jassid in the third week of September with year round occurrence.

2.4 Aphid, *Aphis gossypii*

Aphid belongs to the family Aphididae family and is a major sucking pest of several commercially important food crop. Different species of aphid such as *Aphis craccivora*, *Aphis gossypii*, *Myzus persicae* feed on brinjal, tomato and many other vegetables as well as cereal crops [74]. [75] reported aphid as a serious threat to agricultural crops around the world.

2.4.1 Host preference

Aphid is a versatile crop pest reported from all over the world [76]. Reported 29 plant species of the family Solanaceae to be the host for the *A. gossypii* from India and recognized *C. annuum* as the most preferred one [77]. Prepared a checklist of aphids of Honduras on different host plants and reported *A. gossypii* and *M. persicae* to be the most common on *Solanum melongena*. [78] Reported that *Aphis craccivora* is the most common aphid species infesting a wide range of vegetables and pulse crops.

2.4.2 Nature of damage

[79] Stated that the direct consequences of aphid infestation causes yield losses, decline in quality and increased agricultural potential risks. Aphids can accumulate in high densities on young tender parts of the plants because they have high colonizing capacity; eventually they suck the sap especially from the lower side of the young leaves. Infested plants turn pale, leaves become distorted, curled and crinkled leading to stunted growth of the plants. Aphids secrete honey dew, which encourages growth of sooty molds over the leaves reducing photosynthetic activity of the

crop and can further deter natural enemies of aphids. Severe attack of aphid could lead to production of fruits with reduce size as well as the market value [80].

2.4.3 Succession of aphid in brinjal

Aphid population development in brinjal had a major negative association with maximum temperature, minimum temperature, and rainfall, while relative humidity was positively associated with population size in February with peak aphid population [75]. Studies by [81] showed that *A. gossypii* is abundant in population and spatial distribution and the weather conditions varied. Then, after 56 days of transplantation, max aphid population was found. According to [82] peak aphid nymphal density was reported in March while peak adult aphid abundance was reported in February and March [83]. According to their analysis *A. Gossypii* population started to grow in August and peaked in January and nearly marginal in April at *Solanum melangena* in Bangladesh. In *Solanum* the second period of aphid infestation can be found around May-June [84].

2.5 Whitefly, *Bemisia tabaci*

Whitefly is a polyphagous pest belonging to the family Aleyrodidae of Homoptera order. About 12,000 different species of whitefly were reported worldwide [85]. Importantly, whitefly includes 41 distinctly isolated species population with 24 populations of specific biotypes [86]. Whitefly can cause considerable yield loss and damage to brinjal plants [87].

2.5.1 Host preference

Whitefly is the most abundant and versatile crop pests and known to infest around 600 different crop plants and wild plants [88,89]. Reported that whitefly can attack 500 species of plants belong to 74 taxonomic families. Among the plants squash, tomato, brinjal, potato, pumpkin, cucurbits, okra, beans are noteworthy. Parthenium is one of the most favourite host of whitefly [90]. It also known to feed on weed like Itsit, datura, milkweed, *Chenopodium* sp. [91].

2.5.2 Nature of damage

Whitefly can cause chlorosis, leaf withering, premature leaf drops and wilting of the crop plants by sucking the phloem sap of plant tissue

[92]. Whitefly excretes honeydew which leads to development of sooty mould reducing the effective leaf area for photosynthesis [93]. This may also results in irregular ripening of fruit [94]. Whitefly acts as a vector of virus disease and surprisingly, it can transmit nearly 114 virus species and some can bring havoc to crops [95]. Due to the fact that whitefly lives on the lower side of the leaves, it becomes trouble worthy to control whitefly with contact insecticides [96]. Whitefly is also reported to be possessed resistance to a wide range of insecticides [97].

2.5.3 Succession of Whitefly, *Bamisia tabaci* Genn.

[98] Recorded that white fly activity started from the second week of August (33rd SMW) and lasted until the first week of February. The highest white fly population (19 Nos./plant) was registered in the last week of September (39th SMW), where the maximum and minimum temperature and relative humidity are 34.3, 26.2 and 71.7% accordingly. Whitefly was first recorded in the third week of December (50th SW) according to the experiment of [99] and pest activity continued from the second week of December on until the first week of May. While, he believed that weather factors had no major impact on population dynamics, on the contrary [60] stated that relative humidity had positive effects on the pest population. [100] performed an experiment on the population ecology of whitefly on brinjal and observed peak larval production at 4 to 7 weeks after the crop was transplanted and the highest number was observed on the middle plant canopy. There was no meaningful association between temperature, daylight hours and RH with the populations while wind intensity had a strong correlation. [101] recorded strong plague activity during the third week of February.

3. CONCLUSION

To circumvent the huge production loss and crop damage inflicted by insect pest complex of brinjal, it is important to know the host preferences, nature of damage and seasonal abundance of insect pests for development of an effective management strategy. The pertinent literatures cited and overviews prepared in this paper could help the end users in successful implementation of management tactics of the major insect pests of brinjal in future.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Solanaceae source. A global taxonomic resource for the nightshade family; 2018. Available: www.solanaceaesource.org on 01 July 2020
2. Anonymous. Eggplant. Accessed 25 July 2020. Available: <https://www.tridge.com/intelligences/eggplant/production>
3. Kandoliya UK, Bajaniya VK, Bhadja NK, Bodar NP, Golakiya BA. Antioxidant and nutritional components of eggplant (*Solanum melongena* L.) fruit grown in Saurashtra region. Intl J Curr Microbiol App Sci. 2015;4(2):806-813.
4. Wankhede CB. Studies on pest complex of brinjal with special reference to shoot and fruit borer on different varieties. MS Thesis, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, India; 2009.
5. Rai MK, Pandey AK. Folk medicines of Gond tribe of Seoni district, M.P. India. J Non Timber Forest Prod. 1997;4:61-69.
6. Cao G, Sofic E, Prior RL. Antioxidant capacity of tea and common vegetables. J Agric Food Chem. 1996;44:3426-3431.
7. Magioli C, Mansur E. Eggplant (*Solanum melongena* L.): Tissue culture, genetic transformation and use as an alternative model plant. Acta Botanica Brasilica. 2005;19(1):139-148.
8. Silva ME, Santos RC, O'Leary MC, Santos RS. Effect of aubergene (*Solanum melongena*) on serum and hepatic cholesterol and triglycerides in rats. Braz Arch Biol Technol. 1999;42:339-342.
9. Mutalik S, Paridhavi K, Rao CM, Udupa N. Antipyretic and analgesic effect of leaves of *Solanum melongena* Linn. in rodents. Indian J Pharmacol. 2003;35:312-315.
10. Igwe SA, Akunyili DN, Ogbogu C. Effects of *Solanum melongena* (garden egg) on some visual functions of visually active Igbos of Nigeria. J Ethnopharmacol. 2003;86:135-138.
11. Vohora SB, Kumar I, Khan MS. Effect of alkaloids of *Solanum melongena* on the

- central nervous system. J Ethnopharmacol. 1984;11:331-336.
12. Regupathy A, Palanisamy S, Chandramohan N, Gunathilagaraj K. A guide on crop pests. Sooriya Desk Top Publishers, Coimbatore. 1997;264.
13. Ahmad R. Studies on the pests of brinjal and their control with special reference to fruit borer, *Leucinodes orbonalis* Guen (Pyralidae: Lepidoptera). Entomol News Let. 1974;1(4):2-3.
14. Saimandir J, Gopal M. Evaluation of synthetic and natural insecticides for the management of insect pest control of eggplant (*Solanum melongena* L.) and pesticide residue dissipation pattern. American J PI Sci. 2012;3(2):214- 227.
15. Chakraborty S, Sarkar PK. Management of *Leucinodes orbonalis* Guenee on eggplant during the rainy season in India. J PI Prot Res. 2011;51(4):325-328.
16. Latif MA, Rahman MM, Alam MZ. Efficacy of nine insecticides against shoot and fruit borer, *Leucinodes orbonalis* Guenee (Lepidoptera: Pyralidae) in eggplant. J Pest Sci. 2010;83(4):391-397.
17. Dutta P, Singha AK, Das P, Kalita S. Management of brinjal fruit and shoot borer, *Leucinodes orbonalis* Guenee in agro-ecological condition of West Tripura. Schol J Agric Sci. 2011;1(2):16-19.
18. Thapa RB. Integrated management of brinjal fruit and shoot borer, *Leucinodes orbonalis* Guen: An overview. J Inst Agric Anim Sci. 2010;30:1-16.
19. Choudhury MAR, Rahman MM, Alam MZ, Hossain MM, Khaliq QA. Fluctuation pattern of insect pests and natural enemies in summer brinjal ecosystem. J. Sylhet Agril. Univ. 2015;2(2):149–155.
20. Nayer KK, Krishnan TNA, David BV. General and applied entomology. 11th edn. Tata McGraw- Hill pub. Co. Ltd. 4/12, New Delhi- 110002. 1995;557.
21. Hossain SH, Saqui MAA, Hossain M, Hossain MA, Hossainy NUA, Kabir H. Pest risk analysis (PRA) of brinjal. Plant Quarantine Wing, Department of Agricultural Extension. Dhaka, Bangladesh. 2016;149.
22. Alam MZ, Sana DL. Biology of the brinjal shoot and fruit borer, *Leucinodes orbonalis* G. (Pyralidae: Lepidoptera) in East Pakistan. The Scientist. 1962;5(1-4):13-24.
23. Karim MA. Vegetable and spices insect pests and their control. A lecture in training course on winter vegetable and spice production. Horticulture research and development project. Joydevpur, Gazipur. 1994;75.
24. CABI Crop protection compendium. CAB International; 2007.
(Retrieved on March 15, 2012)
Available:<http://www.cabicompndium.org/cpc>
25. Ishaque NMM, Chaudhuri RP. Larval development behavior of *Leucinodes orbonalis* Guen. reared on some brinjal varieties. J Res Assam Agril Univ. 1984;5:93-97.
26. Onekutu A, Omoloye AA, Odebiyi JA. Biology of the eggfruit and shoot borer (EFSB), *Leucinodes orbonalis* Guenee (Crambidae) on the Garden Egg, *Solanum gilo* Raddi. J Entom. 2013;10(3):156–162.
Available:<https://doi.org/10.3923/je.2013.156.162>
27. Taley YM, Nighut US, Rajurkar BS. Bionomics of brinjal fruit and shoot borer, (*Leucinodes orbonalis* Guenee). Punjabrao Krishi Vidya Res J. 1984;8(1):29-39.
28. Alpuerto AB. Ecological studies and management of brinjal fruit and shoot borer, *Leucinodes orbonalis* Guenee. Indian J Agric Sci. 1994;52(6):391-395.
29. Alam SN, Hossain MI, Rouf FMA, Jhala RC, Patel MG, Rath LK, Sengupta A, Baral K, Shylesha AN, Satpathy S, Shivalingaswamy TM, Cork A, Talekar NS. Implementation and promotion of an IPM strategy for control of eggplant fruit and shoot borer in South Asia. Technical Bulletin. AVRDC publication AVRDC – The World Vegetable Center, Shanhua, Taiwan. 2006;06-672(36):74.
30. Rahman MM. Vegetable IPM in Bangladesh. Department of Entomology. Bangabandhu Sheikh Mujibur Rahman Agricultural University Gazipur, Bangladesh; 2006.
Available:<http://ipmworld.umn.edu/chapters/rahman.htm>
31. Shaukat MA, Mustafa MA, Ammad A, Maqsood S, Umer H, Mustafa F, Malik G. Life aspects and mode of damage of brinjal shoot and fruit borer (*Leucinodes Orbonalis* Guenee) on eggplant (*Solanum Melongena* Linnaeus): A review.

- International J Entom Res. 2018;3(2):28-33.
32. Atwal AS. Agricultural pests of India and Southeast Asia. Kalyani Publishers. New Delhi, India. 1976;529.
33. Baang LA and Corey FM. Life history of an eggplant fruit and shoot borer, *Leucinodes orbonalis* (Guenee) (Lepidoptera: Pyralidae). Chiang Mai Univ J Sci. 1991;4(1):45-61.
34. Islam MN, Karim MA. Management of the brinjal shoot and fruit borer, *Leucinodes orbonalis* Guen. (Lepidoptera: Pyralidae) in field. In: Annual Research Report 1990-91. Entomology Division, Bangladesh Agriculture Research Institute Joydebpur, Gazipur. 1991;44-46.
35. AVRDC. Annual Report. Asian Vegetable Research and Development Centre, Shanhua, Tainan, Taiwan. 1998;148.
36. Butani DK, Jotwani MG. Insects in vegetables. Periodical, Expert Book Agency. New Delhi, India. 1984;356.
37. Neupane FP. Crop pest and their management (4th Edn.) (Nepali language). Sajha Prakashan, Pulchowk, Lalitpur, Nepal. 2001;582.
38. Naqvi AR, Pareek BL, Nanda US, Mitharwal BS. Biophysical characters of brinjal plant governing resistance to shoot and fruit borer, *Leucinodes orbonalis*. Indian J PI Prot. 2009;37(1-2):1-6.
39. Farman A, Inamullah K, Zakirullah AA and Rahman A. Study on the population trend of brinjal fruit borer *Leucinodes Orbonalis* Guen. (Lepidoptera: Pyralidae) and its susceptibility to different insecticides. ARPN J Agric Biol Sci. 2016;11(7):262-266.
40. Ghosh SK, Senapati SK. Seasonal fluctuation in the population of *Leucinodes orbonalis* (Guen.) in the sub-Himalayan region of West Bengal, India and its control on eggplant (*Solanum melongena* L.). Prec Agric. 2009;10(5):443-449.
41. Varma S, Anandhi P, Singh RK. Seasonal incidence and management of brinjal shoot and fruit borer, *Leucinodes orbonalis*. J Entomol Res. 2009;33:323-329.
42. Patel JR, Korat DM, Patel VB. Incidence of shoot and fruit borer (*Leucinodes orbonalis* Guen.) and its effect on yield in brinjal. Indian J PI Prot. 1988;16:143-145.
43. Bhushan S, Chaurasia HK, Shanker R. Efficacy and economics of pest management modules against brinjal shoot and fruit borer (*Leucinodes orbonalis*). The Bioscan. 2011;6(4):639-642.
44. Shukla A, Khatri SN. Incidence and abundance of brinjal shoot and fruit borer *Leucinodes orbonalis* Guenee. The Bioscan. 2010;5(2):305-308.
45. Jamwal VV, Ahmad H, Sharma D. Host biology interactions of *Epilachna vigintioctopunctata* Fabr. The Bioscan. 2013;8(2):513-517.
46. Ahmad M, Ahmad MDJ, Afroze S, Mishra RK. First record of coccinellid beetles (Coleoptera: Coccinellidae) on poplar, *Populus deltoides* from north India. Indian Fores. 2001;127(8):891-897.
47. Wilson RL. Studies of insect feeding on grain amaranth in the Midwest. J Kansas Entomol Soc. 1989;62(4):440-448.
48. Ganga G, Chetty JS. The possible role of reservoir plants in the maintenance of the brinjal pest *Henosepilachna* (*Epilachna*) *vigintioctopunctata* Fabr. (Coleoptera: Coccinellidae) throughout the year. Entomon. 1982;7(2):219-220.
49. Maurice NG, Shukla PK and Ramteke PW. Effect of larval density on survival and occurrence of cannibalism in *Epilachna vigintioctopunctata* Fab. Ann Plant Prot Sci. 2013;21(2):291-294.
50. Imura O, Ninomiya S. Quantitative measurement of leaf area consumption by *Epilachna vigintioctopunctata* (Fabricius) (Coleoptera: Coccinellidae) using image processing. Appl Entomological Zoological. 1978;33:491-495.
51. Srivastava AS, Katiyar SSL. *Epilachna vigintioctopunctata* (Fab.) and *E. dodecastigma* (Muls.) (Coleoptera: Coccinellidae) as a pest of cow pea. Zeits-fur Angew Entomol. 1972;71(2):169-172.
52. Rajagopal D, Trivedi TP. Status, biology and management of epilachna beetle *Epilachna vigintioctopunctata* (Fab.) (Coleoptera: Coccinellidae) on potato in India. Trop. Pest. Manag. 1989;35:410-413.
53. Mall NP, Panday SV, Singh SK. Seasonal abundance of insect pests and estimation of the losses caused by shoot and fruit borer in brinjal. Indian J Entomol. 1992;53:241-247.
54. Jeyasankar A, Premalatha S, Elumalai K. Antifeedant and insecticidal activities of selected plant extracts against epilachna

- beetle, *Epilachna vigintioctopunctata* (Coleoptera: Coccinellidae). Advances Entomol. 2014;2(1):14-19.
55. Omprakash S, Raju SVS. Incidence of *Henosepilachna vigintioctopunctata* Fab. on brinjal in eastern Uttar Pradesh. Intl J Appl Biol Pharma Technol. 2014;5(1):181-183.
56. Haseeb M, Qamar M, Sharma DK. Seasonal abundance of brinjal hadda beetle, *Henosepilachna vigintioctopunctata* F. (Coleoptera: Coccinellidae) in Aligarh, Uttar Pradesh. Trends Biosci. 2009;2(1):31-32.
57. Venkatesha MG. Seasonal occurrence of *Henosepilachna vigintioctopunctata* (F.) (Coleoptera: Coccinellidae) and its parasitoids on Ashwagandha in India. J Asia-Pacific Entomol. 2006;9(3):1-4.
58. Raghuraman M, Veeravel R. Influence of abiotic factors on the incidence of spotted leaf beetle *Henosepilachna vigintioctopunctata* (f.) in brinjal. Pest Manag Hortic Ecosyst. 1999;5:17-20.
59. Ramzan MD, Singh G, Singh GSM, Bhalla JS. Comparative development and seasonal abundance of hadda beetle, *Henosepilachna vigintioctopunctata* (Fabr.) on some solanaceous host plants. J Res Punjab Agric Uni. 1990;27:253-262.
60. Prasad GS, Logiswaran G. Influence of weather factors on population fluctuation of insect pests on brinjal at Madurai, Tamil Nadu. Indian J Entom. 1997;59(4):385 - 388.
61. Das G, Islam T. Relative efficacy of some newer insecticides on the mortality of jassid and whitefly in brinjal. Intl J Res Biol Sci. 2014;4(3):89-93.
62. Ali M, Ashfaq M, Akram W, Sahi ST, Ali A. The physio-morphic characters of the brinjal (*Solanum melongena* L.) plant and their relationship with the jassid (*Amrasca biguttula biguttula* (Ishida)) population fluctuation. Pakistan J Agric Sci. 2012;49(1):67-71.
63. Latif MA, Rahman MM, Islam MR, Nuruddin MM. Survey of arthropod biodiversity in the brinjal field. J Entomol. 2009;6(1):28-34.
64. Iqbal J, Hasan M, Ashfaq M, Sah ST, Ali A. Screening of okra genotypes against jassid, *Amrasca biguttula biguttula* (Ishida) (Homoptera: Cicadellidae). Pak J Agri Sci. 2008;45(4):448-451.
65. Mahmood T, Hussain SI, Khokhar KM, Jeelani G, Mukhtar A. Populations dynamics of leaf hopper (*Amrasca biguttula biguttula*) on brinjal and effects of abiotic factors on it. Asian J Plant Sci. 2002;1(2):107-108.
66. Sharma K, Chander S. Spatial distribution of jassid *Amrasca biguttula biguttula* (Ishida) on cotton. Indian J. Entom. 1998;60(4):326-328.
67. Samal T, Patnaik HP. Field efficacy of insecticides against *Amrasca biguttula biguttula* (Ishida) incidence on eggplant. Ann PI Protec Sci. 2008;16(1):115-118.
68. Deole S. Screening of brinjal cultivars against jassid, *Amrasca biguttula biguttula* based on the leaf texture of the plant. J Appl Zool Res. 2008;19(2):139-140.
69. Sankeshwar M, Patil R. Effect of trichome density on the resistance to the jassids in inter-specific *G. hirsutum* x *G. barbadense* cotton recombinant inbred lines (RIL) population. J Farm Sci. 2016;29(4):479-481.
70. Saroj AK, Singh AK, Singh DC, Kumar P. Population dynamics of jassid *Amrasca biguttula biguttula* (Ishida) in okra, *Abelmoschus esculentus* (L.) and its correlation with meteorological parameters. Plant Arch. 2017;17(2):1189-1190.
71. Gangwar RK, Singh DV. Study on insect pest succession of brinjal crop ecosystem in western region of Uttar Pradesh. India J Biol Agric Health. 2014;4(17):116-119.
72. Dabhi MV, Koshiya DJ. Effect of abiotic factors on population dynamics of leafhopper, *Amrasca biguttula biguttula* (Ishida) in okra. Adv Res J Crop Improv. 2014;5(1):11-14.
73. Dhamdhare S, Dhamdhare SV, Mathur R. Occurrence and succession of pests of brinjal, *Solanum melongena* Linn. at Gwalior (Madhya Pradesh). India J Ent Res. 1995;19(1):71-77.
74. Powell G, Tosh CR, Hardie J. Host plant selection by aphids: Behavioral, Evolutionary, and Applied Perspectives. Ann Rev Entom. 2006;51(1):309-330. Available: <https://doi.org/10.1146/annurev.ento.51.110104.151107>
75. Shakeel M, Akram W, Ali A, Ali MW, Nasim W. Frequency of aphid (*Aphis Gossypii* G.) on brinjal (*Solanum melongena* L.) and farming practices in the agro-climatic

- conditions of Faisalabad, Pakistan Intl J Agric Innov Res. 2014;2(5):841-845.
76. Singh SV, Singh KS, Malik YP. Seasonal abundance and economic losses of shoot and fruit borer, *Leucinodes orbonalis* Guenee on Brinjal. Indian J Entom. 2000;2(5):247-252.
77. Evans GA, Halbert SE. A checklist of the aphids of Honduras (Hemiptera: Aphididae). Florida Entomol. 2007;90:518-523.
78. Nayer KK, Anantakrishna S, Danid BV. General entomology. Mc Grow Hill Publication Co. Ltd. New Delhi, India. 1976;256.
79. Miller GL, Favret C, Carmichael A, Voegtlin DJ. Is there a cryptic species within *Aulacorthum solani* (Hemiptera: Aphididae)? J Econ Entomol. 2009;102:398-400.
80. Ghosh SK, Laskar N, Senapati SK. Seasonal fluctuation of *Aphis gossypii* Glov. on brinjal and field evaluation of some pesticides against *A. gossypii* under Terai region of West Bengal. Indian J Agric Res. 2004;38: 71-177.
81. Touhidur MR, Idris AB, Roff M. Population abundance and spatial distribution of *Aphis gossypii* Glover (Homoptera: Aphididae) and coccinellids on chilli (*Capsicum annum* L.). J Trop Agric Fd Sc. 2006;34:393-403.
82. Rondon SI, Cantliffe DJ, Price JF. Population dynamics of the cotton aphid, *Aphis gossypii* (Homoptera: Aphididae), on strawberries grown under protected structure. Florida Entomol. 2005;88:152-158.
83. Karim KNS, Das BC, Khalequzzaman M. Population dynamics of *Aphis gossypii* Glover (Homoptera: Aphididae) at Rajshahi, Bangladesh. Online J Biological Sci. 2001;1:492-495.
84. Tancik LCJ. Population dynamics of green peach aphid, *Myzus persicae* (Sulz.) (Homoptera, Aphididae) on peach tree (*Persica vulgaris* Mill.) and pepper (*Capsicum annum* L.). Proceedings of the International Scientific Conference on the Occasion of the 55th Anniversary of the Slovak Agricultural University in Nitra. 2001;4:305-307.
85. Bartlett AC, Gawel NJ. Determining whitefly species. Sciences. 1993;261: 1333-1334.
86. Perring TM. The *Bemisia tabaci* species complex. Crop Prot. 2001;20:725-737.
87. Mandal S, Singh NJ, Konar A. Efficacy of synthetic and botanical insecticides against whitefly (*Bemisia tabaci*) and shoot and fruit borer (*Leucinodes orbonalis*) on brinjal *Solanum melongena* L. J. Crop Weed. 2010;6(1):49-51.
88. Cueller ME, Morales FJ. The whitefly *Bemisia tabaci* G. as pest and vector of plant viruses of common bean, *Phaseolus vulgaris* L. Rev. Colombiana Entomol. 2006;32(1):1-9.
89. Arnal E, Russell LM, Debrot E, Ramos F, Carmeli M, Marceno R, Montagne M. A listing of whiteflies and their host plants in Venezuela. Florida Entomol. 1993;76:365-381.
90. Kedar SC, Saini RK, Kumaranag KM. Whitefly, *Bemisia tabaci* (Gennadius) as influenced by host plants in Haryana. Indian J Entom. 2018;80(2):257.
Available:https://doi.org/10.5958/0974-8172.2018.00042.1
91. Muniz M. Host suitability of two biotypes of *Bemisia tabaci* on some common weeds. Entom Exp Appl. 2000;95(1):63-70.
Available:https://doi.org/10.1046/j.1570-7458.2000.00642.x
92. Brown JK. The *Bemisia tabaci* complex: Genetic and phenotypic variability drives *Begomovirus* spread and virus diversification. Online APSnet Features; 2007.
DOI: 10.1094/APSnetFeature-2007-0107
93. Henneberry TJ, Jech LF, de la Torre T. Honeydew production by sweet potato whitefly adult and nymphs. Arizona Cotton Report, The University of Arizona College of Agriculture and Life Sciences. 2001; 37.
94. McKenzie CL, Albano JP. The effect of time of sweet potato whitefly infestation on plant nutrition and development of tomato irregular ripening disorder. Hort Tech. 2009;19(2):353-359.
95. Jones DR. Plant virus transmitted by whiteflies. European J Plant Path. 2003;109:195-219.
96. Zhang W, McAuslane JH, Schuster DJ. Repellency of ginger oil to *Bemisia argentifolii* (Homoptera: Aleyrodidae) on Tomato. J Econ Ent. 2004;97(4):1310-1318.

97. Dennehy TJ, DeGain B, Harpold G, Brown JK, Byrne F, Morin S, Nichols R. First new world report of Q biotype of *Bemisia tabaci* (Gennadius) reveals high levels of resistance to insecticides. Res Pest Manag. News. 2006;15(2):18-20.
98. Sharma N. Studies on pest complex of brinjal with special reference to IPM of brinjal. MS thesis, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior, India; 2012.
99. Ramrao GV. Studies on pest complex of brinjal and their management with microbials. MS Thesis, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, India; 2012.
100. Fauziah I, Fairuz K, Saiful M, Jamaludin M. Population ecology of whitefly, *Bemisia tabaci* (Homoptera: Aleyrodidae) on brinjal. J Agric Sci. 2009;1:27-32.
101. Naik VCB, Rao PA, Krishnayya PV, Chalam MSV. Seasonal incidence and management of *Bemisia tabaci* Gennadius and *Amrasca biguttula biguttula* Ishida of brinjal. Ann PI Protec Sci. 2009;17(1):971-973.

© 2020 Nasif and Siddiquee; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/59627>