Soil Transmitted Helminth Infections among Slum Dwelling Women in Dhaka, Bangladesh

Mandira Mukutmoni^{1*}, Sharmin Musa¹ and Asmaul Hosna²

¹*Department of Zoology, Faculty of Biological Sciences, University of Dhaka, Bangladesh.*

²*Department of Zoology, Faculty of Life and Earth Sciences, Jagannath University, Bangladesh.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors SM and MM designed the study. Author MM wrote the protocol and first draft of the manuscript. Authors MM and AH managed literature searches and statistical analysis. Author MM managed the analysis of the study and author SM edited the last manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2020/v35i830264

Editor(s):

(1) Dr. Gabriel Oladele Awe, Ekiti State University, Nigeria.

Reviewers:

(1) Dirceu Guilherme de Souza Ramos, Federal University of Goiás, Brazil.

(2) José Gabriel Gonçalves Lins, School of Veterinary Medicine and Animal Science - UNESP, Brazil.

(3) Timothy Auta, Federal University Dutsin-Ma, Nigeria.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/59933>

Original Research Article

Received 06 June 2020
Accepted 11 August 2020
Published 19 August 2020

ABSTRACT

Aims: The objective of this trial was to establish the prevalence and risk factors relating to soil-transmitted helminth (STH) infections among slum dwelling women in Dhaka city, Bangladesh.

Study Design: Cross sectional study.

Place and Duration of the Study: A total of 100 stool samples were collected from Ganaktuli (Hazaribagh) slum of Dhaka city from March to September 2019. Women aged between 21 to 40 years old were part of this study, and only those who had at least one primary school going child were included.

Methodology: The interviewed women were asked to provide their stool samples in the containers supplied by the volunteers, and to fill up a questionnaire regarding their socio-demographic and behavioral practices. Formal Ether Concentration technique was applied to process the stool samples, and detection of helminth eggs was done by microscopy.

Results: Of the hundred women, 87 (87%) were infected with at least one STH. *Ascaris lumbricoides* (41.38%) showed the highest prevalence followed by *Trichuris trichiura* (36.78%). Working as day laborers, irregular nail clipping, irregular use of soap after defecation, walking

*Corresponding author: E-mail: mukutmoni.zoo@du.ac.bd;

barefoot, using open and common toilet, irregular consumption of antihelminthic drugs were noticed as significant risk factors.

Conclusion: Soil transmitted helminth infection is still a problem in Bangladesh. Proper drug administration and increase of hygiene practices among the slum dwellers are essential to lessen the high prevalence of soil transmitted helminth infections.

Keywords: Dhaka; slum; women; stool; soil transmitted helminths.

1. INTRODUCTION

Soil-transmitted helminth (STH) infections are considered the most prevalent neglected tropical diseases (NTD) with an estimated number of 1.45 billion people infected with at least one parasite [1,2]. In 2010, 438.9 million people were infected with hookworm, 819.0 million with *Ascaris lumbricoides*, and 464.6 million with *Trichuris trichiura*, globally. Of the 4.98 million years lived with disability (YLDs) referable to STH, 65% were attributable to hookworm, 22% to *A. lumbricoides* and the remaining 13% to *T. trichiura* [2] (The NTDs including STH infection result in prolonged phases of ill health, and in fact support to uphold poverty over their continuing effects on child development and worker productivity [3]. The high STH burden in Asia is probably due to the moist and tropical climatic conditions, scarcity of safe drinking water, inadequate sanitation, and poor hygiene practices, all of which facilitates worm survival and transmission [4,5]. Government of Bangladesh has implemented school-based mass drug administration (MDA) bi-annually since 2008 aimed to control STH infection, administering mebendazole to pre-school children through the Bangladesh Expanded Program on Immunization [6,7]. Regardless of several rounds of MDA, the government is still facing challenges to accomplish the target coverage and utilization of the intervention [6].

Due to urbanization, people tend to come to Dhaka, the capital city of Bangladesh, looking for better employment and income opportunities. A number of immigrants initially concentrate in slums [8]. Within the slums, poor environmental conditions and depressed infrastructures are observed. Due to lack of environmental hygiene, overcrowding, insufficiency of pure water and sanitation, slum dwellers may become affected by various health problems [9]. Moreover, women suffer in the developing countries because of poverty coupled with social and traditional practices, philosophies, gender based ferocity, lack of educations and insufficient healthcare services. Women continue to

experience inferior health outcomes across a number of conditions, despite human rights advances and improvements in certain areas of health and development [10]. The present study was perceived to determine the prevalence of common STH among slum dwelling young women of Dhaka and to assess socio-demographic and behavioral risk factors associated with STH infection among them.

2. MATERIALS AND METHODS

Stool samples were collected from the women residing in Ganaktuli slum which is located in Hazaribagh thana of Dhaka city, Bangladesh.

2.1 Study Population and Design

The present study was conducted among the women aged between 21 to 40 years old and having at least one primary school-going child. A total of 100 stool samples from the 100 women were obtained from March 2019 to September 2019. House to house visits were done by the volunteers. A written questionnaire regarding demographic information and behavioral practices were provided to the women. The volunteers assisted the women who did not know how to read or write. For collecting stool samples, each woman received a properly capped container. They were instructed by the research assistants and the volunteers on how to put an amount of stool into the container. The stool samples were transported to the Parasitology laboratory, Department of Zoology, University of Dhaka within one to two hours of collection into a chiller box. The samples were kept in refrigerator at 4°C and examined within two days of collection.

2.2 Laboratory Screening

Formal Ether Sedimentation Technique was applied to examine the collected samples [11]. The stool samples were emulsified using 4 ml of 10% formal water suspension which was strained to remove large fecal particles. Then 4 ml of

ether was added and the tube mixed for 1 min and immediately centrifuged at 1000 g. After centrifuging, the parasites sedimented to the bottom of the tube and the fecal debris collected in a layer between the ether and formal water. After discarding the supernatant, the sediment was transferred to a slide and covered with a cover glass. The sediment was examined microscopically for cysts, oocysts, eggs and larvae of intestinal parasites. The portion of stool samples were processed and examined microscopically using the 10x objective first. Eggs of helminth parasites were identified according to Chatterjee [12].

2.3 Data Analysis

SPSS version 20.00 was used to input data and to analyze. Chi-square test was applied; the level of significance of each test was set at $P < 0.05$.

3. RESULTS AND DISCUSSION

Among the hundred stool samples examined, 87 samples were positive for at least one STH (Table 1). *A. lumbricoides* (36%) showed the highest prevalence followed by *T. trichiura* (32%). Concurrent infection with *A. lumbricoides* and *T. trichiura* was a little higher (10%) than the hookworm infection (9%) (Table 1). *Ascaris* sp. and *Trichuris* sp. have most often been found in urban and peri-urban communities whereas hookworm is found more often in rural communities [13]. *T. trichiura* is specifically highly prevalent in the tropical and sub-tropical regions and people without proper sanitation are at risk of infection [14]. In 2006, Nguyen et al. conducted a study on reproductive-age Vietnamese women and found that 76% were infected with one or more helminth species; 36% with hookworm, 59% with *A. lumbricoides* and 28% with *T. trichiura* [15]. The egg of *A. lumbricoides* is recognized to stick to dust, fruits and vegetables. Infections usually occur through ingestion of infective ova from contaminated hands, food or drinks. As women usually do the dusting and kitchen chores, especially in underprivileged societies, they possess high risk of STH infections. The high prevalence of ascariasis in our study may be attributed to poor personal hygiene and low economic status of slum dwellers.

Unexpectedly, women having primary education (41.38%) were more vulnerable to STH infection than the women with no institutional education (32.18%) ($P = 0.460$) which is similar to results

found by Suntaravitun and Dokmaikaw [16]. The better educated the parents are, the lower the prevalence of intestinal parasites in children was reported by Nematian et al. [17]. Among the day laborers, prevalence was higher (63.22%) than that of the housewives (36.78%) ($P = 0.006$). Maternal unemployment was considered as risk for infection in children by Quihui et al. [18]. Day laborers have to work outside and they are more exposed to the open environment where hygiene could be less maintained. In our study, all the women using open toilet were STH positive. Women who used common or shared toilet had higher prevalence (50.57%) than the personal toilet users (21.84%) ($P = 0.000$) (Table 2). Though males are more frequently reported open defecating than females [19], we found a high interest among women to not use toilet facilities. Belyhun et al. and Gunawardena et al. found that open defecation increases the risk of hookworm infection [20,21]. Open defecation is likely to be highly contaminated environments with high probability of transmission, predominantly as hookworm larvae can directly enter the skin.

Table 1. Distribution of STH among STH-positive women in Ganaktuli slum of Dhaka city, Bangladesh (n = 87)

Infection	n (%)
<i>A. lumbricoides</i>	36 (41.38)
<i>T. trichiura</i>	32 (36.78)
Hookworm	9 (10.34)
<i>A. lumbricoides</i> and <i>T. trichiura</i>	10 (11.49)
Total	87 (100)

All the women who never used antihelminthic drugs in lifetime, were STH positive. The women who used to take antihelminthic drugs once in year or two years were categorized as irregular users and the women who used to take at least twice in year were categorized as regular users. Both the irregular and regular users, used to take both albendazole and mebendazole. Irregular antihelminthic drug consumption resulted in higher STH prevalence (44.83%) than using at regular time interval (33.33%) ($P = 0.008$). Northrop-Clewes et al. found that treatment with mebendazole reduces the prevalence of *A. lumbricoides* from 78% to 8%, of *T. trichiura* from 65% to 9% and of hookworm from 4% to 0% [22]. According to Hall and Nahar, albendazole was found to act mildly against *A. lumbricoides* but within 10 days, all dosages had cured about 92% of infection. For the reduction of *T. trichiura* egg,

they found 400 mg of albendazole for 3 days to achieve a cure rate of 80% [23].

We observed, 34 women used to clip their nails irregularly (only once in three to five months) and among them 33 (37.93%) were STH positive ($P = 0.032$) (Table 2). Mahmud et al. conducted a study among the children of northern Ethiopia, and they revealed that whose nails were cut on a weekly basis were 49% less likely to be re-infected by intestinal parasites than children not receiving the nail clipping intervention [24]. We observed that the women who did not wash their hands with soap after defecation, were all STH positive ($P = 0.004$). Infrequent use of soap was

more likely to maternal soil transmitted helminth infections in Butajira, Ethiopia [20]. In 2019, Gebrehiwet et al. found that the women who had no habit of using soap after any procedure were five times more likely to have the soil transmitted helminth infections compared to those who had practice of washing hands using water and soap [25]. Appropriate handwashing with soap and weekly trimming of fingernails can reduce the output of infective stages in feces that results in the contamination of the environment and therefore can lessen the transmission in the community [26,27]. A review found inconclusive confirmation that hand-washing can lessen *A. lumbricoides* infection [28].

Table 2. Demographic profile of the women residing in Ganaktuli slum of Dhaka city, Bangladesh in relation to STH infection

Demographic profile	STH Positive (n=87), n%	STH negative (n=13), n%	Total (n=100), n%	χ^2 value, df, P-value
Education				
No institutional education	28 (32.18)	2 (15.38)	30 (30.00)	1.551, 2, 0.460
Primary	36 (41.38)	7 (53.85)	43 (43.00)	
Secondary	23 (26.44)	4 (30.77)	27 (27.00)	
Occupation				
Housewife	32 (36.78)	10 (76.92)	42 (42.00)	7.481, 1, 0.006*
Day laborer	55 (63.22)	3 (23.08)	58 (58.00)	
Toilet				
Open	24 (27.59)	0	24 (24.00)	21.488, 2, 0.000*
Common or shared	44 (50.57)	2 (15.38)	46 (46.00)	
Personal	19 (21.84)	11 (84.61)	30 (30.00)	
Anthelmintic drugs				
Never	19 (21.84)	0	19 (19.00)	9.623, 2, 0.008*
Irregular	39 (44.83)	3 (23.08)	42 (42.00)	
Regular	29 (33.33)	10 (76.92)	39 (39.00)	
Nail clipping				
Irregular	33 (37.93)	1 (7.69)	34 (34.00)	4.609, 1, 0.032*
Regular	54 (62.07)	12 (92.31)	66 (66.00)	
Use of shoes				
Irregular	42 (48.27)	2 (15.38)	44 (44.00)	4.966, 1, 0.026*
Regular	45 (51.72)	11 (84.61)	56 (56.00)	
Handwash after defecation				
Without soap	36 (41.38)	0	36 (36.00)	8.406, 1, 0.004*
With soap	51 (58.62)	13 (100)	64 (64.00)	
Keeping animal				
No animal	51 (58.62)	9 (69.23)	60 (60.00)	0.531, 1, 0.466
Keep animal	36 (41.38)	4 (30.77)	40 (40.00)	
Knows about STH				
Never heard	38 (43.68)	6 (46.75)	44 (44.00)	0.028, 1, 0.867
Heard about IP	49 (56.32)	7 (53.85)	56 (56.00)	
Instruct child to wash hand after defecating				
Yes	48 (55.17)	8 (61.54)	56 (56.00)	0.186, 1, 0.666
No	39 (44.83)	5 (38.46)	44 (44.00)	

[χ^2 = Chi-squared value, df=degree of freedom, P value *significant]

In our study, less frequent use of shoes resulted in higher prevalence (48.27%) than the regular use (51.72%) ($P = 0.026$) (Table 2). Gebrehiwet et al. observed that the women who used to wear shoes were 95% less likely to be infected by soil transmitted helminths than who did not wear shoes [25]. Kaliappan et al. reported that poor practice of footwear does not significantly enhance the risk of STH infection [29]. Animal keeping did not show significant influence in STH prevalence in our study ($P = 0.466$) (Table 2). Among the 100 interviewed women, 44 women never heard of STH and among them 38 (43.68%) were STH positive, and 56 women were cognizant of STH; among them 49 (56.32%) were STH positive (Table 2). We observed that 48 (55.17%) STH infected women instructed their children to wash hands with soap after defecating and 39 women (44.83%) did not use to instruct (Table 2). Kassaw et al. observed that 45.20% of the women knew about and 54.8% of women did not know about intestinal parasitic infestations, prevention and control methods in their study in Sekota town [30]. According to the findings of Kamunvi and Ferguson, intestinal worms rank poorly in people's minds as a vital health problem [31].

4. CONCLUSION

The study illustrated high prevalence of soil transmitted helminths among the slum dwelling women. Improper use of antihelminthic drugs, bad practice of defecation in open areas and not washing hands properly after defecation are vital issues of concern which should be addressed properly. Consequently, public awareness about STH prevention and control should be generated through campaigns and well-organized trainings.

CONSENT AND ETHICAL APPROVAL

Prior to the commencement of the project entitled "Knowledge, attitude and practice (KAP) assessment of intestinal parasitic and urinary tract infections (UTI) among women and school children in slum areas of Dhaka", ethical clearance was given by the Ethical Review Committee of Faculty of Biological Sciences, University of Dhaka. The volunteers who obtained stool samples from the interviewed, were clearly informed on the aims and objectives of the study. Bengali language (as native) was used during data collection for better communication. As per international standard or university standard, respondents' written consent has been collected and preserved by the authors.

ACKNOWLEDGEMENT

The research was funded by University Grants Commission of Bangladesh. We are thankful for the support.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hotez PJ, Alibek K. Central Asia's hidden burden of neglected tropical diseases. *PLoS Negl. Trop. Dis.* 2011;5(9):e1224. DOI: 10.1371/journal.pntd.0001224
2. Pullan RL, Smith JL, Jasrasaria R, Brooker SJ. Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasit. Vectors.* 2014; 7:37. PMID: 24447578 DOI: 10.1186/1756-3305-7-37
3. Hotez PJ, Fenwick A, Savioli L, Molyneux DH. Rescuing the bottom billion through control of neglected tropical diseases. *Lancet.* 2009;373(9674):1570–1575. PMID: 19410718 DOI: 10.1016/S0140-6736(09)60233-6
4. Brooker S, Singhasivanon P, Waikagul J, Supavej S, Kojima S, Takeuchi T, et al. Mapping soil-transmitted helminths in Southeast Asia and implications for parasite control. *Southeast Asian J. Trop. Med. Public Health.* 2003;34(1):24–36. PMID: 12971511
5. Strunz EC, Addiss DG, Stocks ME, Ogden S, Utzinger J, Freeman MC. Water, sanitation, hygiene, and soil-transmitted helminth infection: A systematic review and meta-analysis. *PLoS Med.* 2014;11(3): e1001620. PMID: 24667810
6. Nath TC, Padmawati RS, Murhandarwati EH. Barriers and gaps in utilization and coverage of mass drug administration program against soil-transmitted helminth infection in Bangladesh: An implementation research. *J. Infect. Public Health.* 2019;12:205-212. DOI: 10.1016/j.jiph.2018.10.002
7. Benjamin-Chung J, Nazneen A, Halder AK, Haque R, Siddique A, Uddin MS, et al. The interaction of deworming, improved sanitation, and household flooring with soil-transmitted helminth infection in rural

- Bangladesh. PLoS Negl. Trop. Dis. 2015;9(12):e0004256.
DOI: 10.1371/journal.pntd.0004256
8. Angeles G, Lance P, Barden-O'Fallon J, Islam N, Mahbub AQM, Nazem N. The 2005 census and mapping of slums in Bangladesh: Design, select results and application. Int. J. Health Geogr. 2009;8: 32-10.
DOI: 1186/1476-072X-8-32
9. Sclar ED, Garau P, Carolini G. The 21st century health challenge of slums and cities. Lancet. 2005;365:901-903.
PMID: 15752535
DOI: 10.1016/S0140-6736(05)71049-7
10. Davidson PM, McGrath SJ, Meleis AI, Stern P, Digiacomo M, Dharmendra T, et al. The health of women and girls determines the health and well-being of our modern world: A white paper from the International Council on women's health issues. Health Care Women Int. 2011; 32(10):870-886.
PMID: 21919625
DOI: 10.1080/07399332.2011.603872
11. Cheesbrough M. Medical laboratory manual for tropical countries II. Cambridge University, New York; 1987.
12. Chatterjee KD. In: Parasitology (Protozoology and Helminthology), 12th edn, Chatterjee Medical Publishers, Calcutta; 1980.
13. Pullan RL, Brooker SJ. The global limits and population at risk of soil-transmitted helminth infections in 2010. Parasit Vectors. 2012;5:81.
PMID: 22537799
DOI: 10.1186/1756-3305-5-81
14. Guerrant RL, Walker DH, Weller PF. Tropical infectious diseases: Principles, pathogens and practice. 3rd Edn. Saunders/Elsevier. Edinburgh; 2011.
15. Nguyen PH, Nguyen KC, Nguyen TD, Le MB, Bern C, Flores R, Martorell R. Intestinal helminth infections among reproductive age women in Vietnam: Prevalence, co-infection and risk factors. Southeast Asian J. Trop. Med. Public Health. 2006;37(5):865-874.
PMID: 17333727
16. Suntaravitun P, Dokmaikaw A. Prevalence of intestinal parasites and associated risk factors for infection among rural communities of Chachoengsao Province, Thailand. Korean J. Parasitol. 2018;56(1): 33-39.
PMID: 29529848
17. Nematian J, Nematian E, Gholamrezanezhad A, Asgari AA. Prevalence of intestinal parasitic infections and their relation with socio-economic factors and hygienic habits in Tehran primary school students. Acta Trop. 2004;92(3):179-86.
PMID: 15533285
DOI: 10.1016/j.actatropica.2004.06.010
18. Quihui L, Valencia ME, Crompton DW, Phillips S, Hagan P, Morales G, Díaz-Camacho SP. Role of the employment status and education of mothers in the prevalence of intestinal parasitic infections in Mexican rural schoolchildren. BMC Public Health. 2006; 6:225.
PMID: 16956417
DOI: 10.1186/1471-2458-6-225
19. Greenland K, Dixon R, Khan SA, Gunawardena K, Kihara JH, Smith JL, et al. The Epidemiology of soil-transmitted helminths in Bihar State, India. PLoS Negl. Trop. Dis. 2015;9(5):e0003790.
DOI: 10.1371/journal.pntd.0003790
20. Belyhun Y, Medhin G, Amberbir A, Erko B, Hanlon C, Alem A, et al. Prevalence and risk factors for soil transmitted helminth infection in mothers and their infants in Butajira, Ethiopia: A population based study. BMC Public Health. 2010;10:21.
PMID: 20085635
DOI: 10.1186/1471-2458-10-21
21. Gunawardena K, Kumarendran B, Ebenezer R, Gunasingha MS, Pathmeswaran A, de Silva N. Soil transmitted helminth infections among plantation sector schoolchildren in Sri Lanka: prevalence after ten years of preventive chemotherapy. PLoS Negl. Trop. Dis. 2011;5(9):e1341.
PMID: 21980549
DOI: 10.1371/journal.pntd.0001341
22. Northrop-Clewes CA, Rousham EK, Mascie-Taylor CGN, Lunn PG. Anthelmintic treatment of rural Bangladeshi children: Effect on host physiology, growth, and biochemical status. American J. Clin. Nutr. 2001;73(1):53-60.
DOI: 10.1093/ajcn/73.1.53
23. Hall A, Nahar Q. Albendazole and infections with *Ascaris lumbricoides* and *Trichuris trichiura* in children in Bangladesh. Trans. R. Soc. Trop. Med. Hyg. 1994;88(1):110-112.
DOI: 10.1016/0035-9203(94)90525-8

24. Mahmud MA, Spigt M, Bezabih AM, Pavon IL, Dinant GJ, Velasco RB. Efficacy of handwashing with soap and nail clipping on intestinal parasitic infections in school-aged children: A factorial cluster randomized controlled trial. *PLoS Med.* 2015;12(6):e1001837.
PMID: 26057703
DOI:10.1371/journal.pmed.1001837
25. Gebrehiwet MG, Medhaniye AA, Alema HB. 2019. Prevalence and associated factors of soil transmitted helminthes among pregnant women attending antenatal care in Maytsebri primary hospital, North Ethiopia. *BMC Res. Notes.* 2019;12:644.
DOI: 10.1186/s13104-019-4684-3
26. Harhay MO, Horton J, Olliaro PL. Epidemiology and control of human gastrointestinal parasites in children. *Expert Rev. Anti. Infect. Ther.* 2010;8:219–234.
PMID: 20109051
DOI: 10.1586/eri.09.119
27. Anderson RM, Truscott JE, Pullan RL, Brooker SJ, Hollingsworth TD. How effective is school-based deworming for the community-wide control of soil-transmitted helminths? *PLoS Negl. Trop. Dis.* 2013;7:e2027.
PMID: 23469293
DOI: 10.1371/journal.pntd.0002027
28. Fung IC, Cairncross S. Ascariasis and handwashing. *Trans. R. Soc. Trop. Med. Hyg.* 2009;103(3):215–22.
PMID: 18789465
DOI: 10.1016/j.trstmh.2008.08.003
29. Kaliappan SP, George S, Francis MR, Kattula D, Sarkar R, Minz S, et al. Prevalence and clustering of soil-transmitted helminth infections in a tribal area in southern India. *Trop. Med. Int. Health.* 2013;18:1452–62.
PMID: 24237860
DOI: 10.1111/tmi.12205
30. Kassaw MW, Abebe AM, Abate BB, Zemariam AB, Kassie AM. Knowledge, attitude and practice of mothers on prevention and control of intestinal parasitic infestations in Sekota Town, Waghimra Zone, Ethiopia. *Pediatric Health Med. Ther.* 2020;11:161-169.
PMID: 32607049
DOI: 10.2147/PHMT.S229610
31. Kamunvi F, Ferguson AG. Knowledge, attitudes and practices (KAP) of human intestinal helminths (worms) in two rural communities in Nyanza Province, Western Kenya. *East Afr. Med. J.* 1993;70(8):482-490.
PMID: 8261967

© 2020 Mukutmoni et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/59933>